

THE CHORD CONSTRUCTION

DAVID LEHAVI AND RON LIVNÉ

ABSTRACT. Let F be a smooth plane curve of degree 3. Let β be an element in $\text{Pic}(F)[2] - \{0\}$. Let us define

$$F' := \{\overline{p(p+\beta)} \mid p \in F\} \subset (\mathbb{P}^2)^*.$$

In this note we show that F' is a smooth embedding of F/β . Moreover, let β' be the generator of $\text{Pic}(F)/\beta$, and let $p \in F$ be a flex, then $\overline{p(p+\beta)} + \beta'$ is a flex on F' . We present two proofs.

1. Notation. Let F be a smooth curve of genus 1, $\beta \in \text{Pic}(F)[2]$. Let us denote $F' := F/\beta$. Let π be the quotient map $F \rightarrow F'$ and let β' be the image of the non-zero element in $\text{Pic}(F)[2]/\beta$ in $\text{Pic}(F')$. Let H be an element in $\text{Pic}^3(F)$, and $i : F \rightarrow \mathbb{P}^2$ the map associated to H . This notation will hold in the next definition and couple of lemmas.

2. Definition (The chord construction (c.c.)). Let us define

$$\begin{aligned} i' : F/\beta &\rightarrow \mathbb{P}^{2*} \\ \{p, p+\beta\} &\mapsto \overline{i(p)i(p+\beta)} \end{aligned}$$

3. Theorem. *The c.c is an embedding.*

Proof. Let us identify F with its image $i(F)$.

The c.c. is 1-1: Let H be the line corresponding to a point $\{p, \beta+p\} \in F/\beta$. Then $F \cap H$ consists of three points:

$$\{p, p+\beta, H - (2p+\beta)\}.$$

If $(H - (2p+\beta)) - \beta \neq p, p+\beta$, then $i'(H)$ consist of the unique point $\{p, p+\beta\}$, so it is 1-1. If $(H - (2p+\beta)) - \beta = p$ then $H - (2p+\beta) = p+\beta$, so i' is 1-1 here too. The remaining case is similar. (in the last two cases p (respectively $p+\beta$) is a flex). For $p \in \mathbb{P}^2$, let p^* be the line it defines in $\mathbb{P}^{2*}$. *If F does not have a CM then for a generic $p \in F$, p^* is transversal to F' :* Suppose it is not. By upper-semicontinuity, it is never transversal, so F is the dual curve of $i'(F')$. Since $i'(F')$ is not a line (it is an immersion of F'), this induces a degree 1 map $i'(F') \rightarrow F$, and therefor a degree 1 map $F' \rightarrow F$. This means that F has a CM.

Date: February 1, 2008.

1991 Mathematics Subject Classification. 14N15, 14H52.

Key words and phrases. Plane cubics, Chords.

The c.c is of degree 3: Assume first that F does not have a CM. Choose a generic $p \in F$. The degree of i' is the number of lines through p which belong to $i'(F')$. There are exactly two types of such lines:

- The line is $\overline{p(p + \beta)}$.
- The line is the one passing through a pair of the four solutions of $x + (x + \beta) = H - p$.

Now suppose F has a CM. Then F and the c.c. are a flat limit of curves without a CM, so the degree of i' is constant. (see [Ha] preposition III.9.8) $\square$

4. Theorem. *If $i(p)$ is a flex, then so is $i'(\pi(p) + \beta')$.*

Proof. We are going to drop the i, i' notation, and identify both curves with their images.

Involutions on $F \subset \mathbb{P}^2$: Let p be a flex on F . Let α be the unique involution of $\mathbb{P}^2$ that

- sends F to itself.
- fixes p .

Note that this involution has a fixed line L , where $L \cap F$ consists of the translates of p by points in $\text{Pic}(F)[2] - \{0\}$. All the involutions of $\mathbb{P}^2$ that send F to itself are of this form (they have to send flexes to flexes, and there are 9 flexes, so one of them is fixed). The same holds for F' .

The dual involution: Given α , there is a dual involution α' on $\mathbb{P}^{2*}$. By the definition of the c.c, α' sends F' to itself. The fixed line and point of α' are p^*, L^* respectively. There is exactly one pair of points in F with difference β on L . This is the pair $\{p + \gamma, p + \gamma + \beta\}$ where γ is some element in $\text{Pic}(F)[2] - \{0, \beta\}$. $\square$

5. We develop a formula for the c.c. . Theorem 10 (which states the formula) also gives an alternative proof for theorems 3,4.

6. Notation. Let us denote homogeneous coordinates on $\mathbb{P}^2$ by $[X; Y; Z]$. Let us denote the dual coordinates on $\mathbb{P}^{2*}$ by $[U; V; W]$. Let us also define

$$x := X/Z, y := Y/Z,$$

and denote the $\mathbb{A}^2$ coordinates by (x, y) Let F as the closure of the locus of $y^2 = f$ where

$$f(x) = x^3 + ax^2 + bx.$$

Let us identify F with $\text{Pic}(F)$ by choosing $O := [0; 1; 0]$ be the zero. Let β be the point $(0, 0) = [0; 0; 1]$.

7. Lemma. *Let $p = (x(p), y(p))$ be a point on F . Then*

$$p + \beta = (bx(p), -by(p)/x(p)^2)$$

Proof. There are exactly three points in

$$F \cap \overline{\beta p},$$

these are $p, \beta, -p - \beta$. Since $\overline{\beta p}$ is given by the nulls of $y/x = \frac{y(p)}{x(p)}$, They are the solutions of

$$(x \frac{y(p)}{x(p)})^2 = x^3 + ax^2 + bx.$$

One of the solutions of this equation is 0. the other two are the roots of

$$0 = x^2 + (a - (\frac{y(p)}{x(p)})^2)x + b.$$

Since one of the roots is $x(p)$, we see that the other is

$$b/x(p) = x(p + \beta).$$

We also deduce that

$$y(p + \beta) = -y(-p - \beta) = -\frac{y(p)}{x(p)}b/x(p) = -by(p)/x(p)^2.$$

□

8. Lemma. *The lines of the c.c. are given in $\mathbb{P}^{2*}$ by*

$$[y(p)(x(p)^2 + b); bx(p) - x(p)^3; -2bx(p)y(p)],$$

where p are the points on F .

Proof. Suppose the line $\overline{p(\beta + p)}$ is given by

$$(1) \quad sx + c = y.$$

In this case we have:

$$s = \frac{y(p) - y(p + \beta)}{x(p) - x(p + \beta)} = \frac{y(p) + by(p)/x(p)^2}{x(p) - b/x(p)} = \frac{y(p)(x(p)^2 + b)}{x(p)^3 - bx(p)}$$

$$C = y(p) - sx(p)$$

Simplifying

$$[s; -1; c],$$

We get the result. □

9. Lemma. *Assuming the c.c. is of degree 3, it satisfies the equation*

$$e(U - \mu^{-1}W)V^2 = W^3 - \frac{2\mu b}{\mu^2 - b}(U - \mu^{-1}W)W^2 - \frac{\mu^2 b}{\mu^2 - b}(U - \mu^{-1}W)^2W.$$

for some e , where $\mu = 2b/a$.

Proof. Let us consider the involution $(x, y) \mapsto (x, -y)$ on $\mathbb{P}^2$. It has a fixed line l : the nulls of $y = 0$, and a unique fixed point out of l : O . If we consider the dual involution, we see that it has a fixed line at the dual of O , to be denoted O^* . It also has a unique fixed point out of O^* . This is the point dual to l , to be denoted l^* . This means that l^* is a flex of the c.c. and its translations by elements of order 2, all sit on O^* . As in the proof of lemma 3, there are two types of such lines:

- The line is $\overline{O\beta}$.
- The line is the one passing through a pair of the four solutions of $2x = \beta$.

In the first type, this is the line $x = 0$. In the second type, we are looking for the solutions of

$$\begin{aligned} x \neq 0, y^2 = f(x), x \frac{dy}{dx} = y &\iff \\ x \neq 0, x \frac{df}{dx} = f &\iff \\ 3x^2 + 2ax + b = 2x^2 + 2ax + 2b &\iff \\ x^2 = b. \end{aligned}$$

We have to identify the tangent to the flex. A tangent to the flex (on a curve of degree 3) is a line that intersect the curve only in the flex. In dual coordinates: It is the unique point $q = (\mu, 0)$ on the line $y = 0$, such that $y = 0$, is the ONLY c.c line on which q sits. The x coordinates of the intersection between a chord and $y = 0$, can be obtained from equation 1, letting $y = 0$. i.e. they are:

$$-c/s = \frac{y(p) - sx(p)}{s} = y(p)/s - x(p) = \frac{x(p)^3 - bx(p)}{x(p)^2 + b} - x(p) = \frac{-2bx(p)}{x(p)^2 + b}.$$

We see that each such point is given by only two $x(p)$. In q , these $x(p)$ are given by the roots of

$$x^2 + ax + b.$$

Whence,

$$\mu = 2b/a.$$

Let us apply the projective transformation

$$U' = U - 1/\mu W, V' = V, W' = W,$$

and define

$$w' = W'/U', v' = V'/U'.$$

In these coordinates we have

$$\begin{aligned} ev'^2 &= w'(w' - \mu\sqrt{b}/(\mu - \sqrt{b}))(w' + \mu\sqrt{b}/(\mu + \sqrt{b})) \\ &= w'(w'^2 - \frac{2\mu b}{\mu^2 - b}w' - \frac{\mu^2 b}{\mu^2 - b}) \end{aligned}$$

for some e . Going back to our original coordinates, we get

$$e(U - \mu^{-1}W)V^2 = W^3 - \frac{2\mu b}{\mu^2 - b}(U - \mu^{-1}W)W^2 - \frac{\mu^2 b}{\mu^2 - b}(U - \mu^{-1}W)^2 W.$$

□

10. Theorem. *The c.c is of degree 3, and $e = \frac{-8b^3}{a^2 - 4b}$*

Proof. This is a trivial check, best done by a computer. One has to define:

$$U := y(x^2 + b), V := bx - x^3, W := -2bxy, \mu := 2b/a,$$

and verify that

$$\frac{W^3 - \frac{2\mu b}{\mu^2 - b}(U - \mu^{-1}W)W^2 - \frac{\mu^2 b}{\mu^2 - b}(U - \mu^{-1}W)^2 W}{(U - \mu^{-1}W)V^2} = -\frac{8b^3 y^2}{(a^2 - 4b)f(x)}$$

□

11. Remark. If F is as before, and $\beta \in \text{Pic}^0(F) - \text{Pic}^0(F)[2]$, the chord construction gives a map of degree 1. The image of this map is of degree 6.

REFERENCES

[Ha] R. Hartshorne *Algebraic Geometry* Springer Verlag, GTM 52. (1977)

MATHEMATICS INSTITUTE, THE HEBREW UNIVERSITY OF JERUSALEM, GIVAT RAM,
JERUSALEM 91904, ISRAEL

E-mail address: dlehavi@math.huji.ac.il

MATHEMATICS INSTITUTE, THE HEBREW UNIVERSITY OF JERUSALEM, GIVAT RAM,
JERUSALEM 91904, ISRAEL

E-mail address: rlivne@math.huji.ac.il